

US009550985B2

## (12) United States Patent

## Tan et al.

# (10) Patent No.: US 9,550,985 B2

(45) **Date of Patent: Jan. 24, 2017** 

## (54) METHODS FOR FORENSIC DNA QUANTITATION

(75) Inventors: Eugene Tan, Arlington, MA (US); Richard F Selden, Lincoln, MA (US);

Rosemary S. Turingan, Stoneham, MA

(US)

(73) Assignee: NetBio, Inc., Waltham, MA (US)

(\*) Notice: Subject to any disclaimer, the term of this

patent is extended or adjusted under 35

U.S.C. 154(b) by 0 days.

(21) Appl. No.: 12/816,370

(22) Filed: Jun. 15, 2010

(65) **Prior Publication Data** 

US 2011/0008785 A1 Jan. 13, 2011

## Related U.S. Application Data

- (60) Provisional application No. 61/268,770, filed on Jun. 15, 2009.
- (51) Int. Cl. C12Q 1/68 (2006.01) G01N 27/447 (2006.01) C12N 15/10 (2006.01) B01L 3/00 (2006.01) B01L 7/00 (2006.01)
- (58) Field of Classification Search

None

See application file for complete search history.

## (56) References Cited

#### U.S. PATENT DOCUMENTS

#### FOREIGN PATENT DOCUMENTS

EP 1 026 258 A2 8/2000 EP 1 614 466 A2 1/2006 (Continued)

## OTHER PUBLICATIONS

Schmalzing et al. PNAS. 1997. 94: 10273-10278.\* (Continued)

Primary Examiner — Joseph G Dauner (74) Attorney, Agent, or Firm — Proskauer Rose LLP

#### (57) ABSTRACT

Described herein are methods and devices for nucleic acid quantification and, in particular, to microfluidic methods and devices for nucleic acid quantification. In certain embodiments methods of quantifying a target nucleic acid without the need for amplification are provided. The methods involve, in some embodiments, allowing a binding agent to become immobilized with respect to the target nucleic acid. In some cases, the binding agent comprises a signaling moiety that can be used to quantify the amount of target nucleic acid. In another aspect, the quantification can be carried out rapidly. For example, in certain embodiments, the quantification can be completed within 5 minutes. In yet another aspect, samples containing a low amount of target nucleic acid can be quantified. For instance, in some cases, samples containing less than 100 nanograms per microliter may be quantified. Also described are devices and kits for performing such methods, or the like.

## 90 Claims, 27 Drawing Sheets

